

Biological Nitrogen fixation

Nitrogen is the greatest limiting factor after water for plant growth. In nature fixed nitrogen is present in organic form as part of proteins and in many inorganic form as NH_3 , NO_3^- . In atmosphere there is abundance of free molecular nitrogen.

The molecular nitrogen can not be directly used by the plant. It is either fixed industrially or biologically by a free living micro-organism. About 95% of the natural fixation of nitrogen occurs through certain micro-organism i.e. Biological nitrogen fixation.

The process of nitrogen transformation is as follows:-

1. Nitrogen fixation: The micro-organism remains in free living state or in symbiotic association.

Non-Symbiotic bacteria :-

These bacteria live freely in the soil and fix atmospheric nitrogen from the air. The nitrogen combines with other element forming nitrogenous compounds. The common non-symbiotic bacteria are Azotobacter (aerobic form) and Clostridium (non-aerobic form).


Symbiotic nitrogen fixing bacteria :-

There are several genera of symbiotic nitrogen fixing bacteria of which Rhizobium is most common. Rhizobium is a gram-negative bacteria which forms association with the members of Fabaceae e.g. Pea, Bean, Medicago etc. Recently, rhizobium has been reported as


The nodules of *Trifolium asperum*

The nodules or the tubercles the home of million of bacteria have the ability to take up free nitrogen of the air and convert it into nitrogenous compounds. A part of the fixed nitrogen passes into the tissues of legume plants and a part diffuses into the surrounding soil.

During infection of a root free living Rhizobium aggregate around root hairs. They penetrate by means of infection thread. Coincident with the passage of the thread, cortical cell division is stimulated to form the bulk of the eventual nodule tissue of polyploid cells. The infection thread usually terminates in a polyploid host cell. Many small groups bacteroids are also formed in the nodule cells. The bacteria utilize the carbohydrate and other food of leguminous plants. The bacteria utilize atmospheric nitrogen to synthesize organic, nitrogenous compounds. Some of these nitrogenous compounds become available to the leguminous plant in which the bacteria is growing.



Infection thread formed by bacteria



Penetration of rhizobia and formation of nodules.


Leghaemoglobin & mechanism of symbiotic N₂ fixation in nodule.

The core of nodule is red due to the pigment, leghaemoglobin - a product of rhizobium-legume complex. Leghaemoglobin is an oxygen carrier, the oxygen is necessary for the electron transport chain of the rhizobium bacteroid. It provides oxygen to the root nodule bacteria quickly, even at very low levels of free oxygen. It is believed that the leghaemo-



glabin levels of molecular oxygen low in the bacteroid. This function is particularly important because nitrogenase is sensitive to O_2 and loses activity in its presence. Nodules lacking leghaemoglobin may be unable to fix nitrogen.

The reduction of nitrogen to ammonia is catalyzed by the enzyme nitrogenase. Certain micronutrients such as iron, copper, cobalt and molybdenum appears to be essential. The first three micronutrients are required or contained in leghaemoglobin synthesis whereas molybdenum functions alternately as an electron acceptor and donor in the reduction of nitrogen to ammonia.

N_2 is reduced to diimide ($H-N=N-H$), hydrazine (H_2N-NH_2) and then ammonia (NH_3) which may be outlined as follows:



OR



Nogales & Fritz



The reduction of nitrogen

into ammonia by the enzyme nitrogenase in bacteroids depend upon availability of ATP and reduced substrate. Capable of elongating hydrogen atoms to nitrogen. ATP is generated in bacteroid respiratory chain system and reduced substrate is obtained from host - cells. Glucose 6 phosphate is considered, to be the reduced substrate for the process and reduced NADP together with ferredoxin function as electron carriers.

Animals and plant proteins are decomposed by bacteria and amino acids are released which gives ammonia by ammonifying bacteria.

Nitrogen converts in Soil


Oxidation of ammonia to nitrate in the soil may occur through autotrophic bacteria Nitrosomonas and Nitrobacter as follows —



The conversion of ammonia to nitrite and then to nitrate is called nitrification.

In denitrification, nitrate

is converted to nitrous oxide and then to N_2 gas. The cycle can be shown as follows —



The blue green algae are also nitrogen fixers. Perhaps, in not too distant future, oriented cropping systems will be made up of culture of nitrogen-fixing algae and one crop plants.

Scientists are busy to transfer the nitrogen fixing gene (*nif*-gene) — unique inherent property of nitrogen fixation in certain microbes to the cereals, vegetables, fruit plants in order to enable the latter, to meet their assimilable N requirement by their own.