

S.B.Sc Part II (Physics Home) ?

S.B College, Agra

Dr. S. D. Singh (Dr Satyadev Narayan Singh)

Q. What do you mean by optical path. State and explain Fermat's principle of extremum path and use it to the law of reflections and refraction of light?

Ans

If a ray of light travels a distance  $d'$  in a medium of refractive index  $n$  than the product  $nd$  is known as optical path in the medium.

Optical path = velocity of light in air  $\times$  time of travel

Fermat's Principle :-

In 1658 Fermat state the principle of extremum path known after him. It is as follows if a ray of light travelling from one point to another by any number of reflections and refractions follows the particular path for which the time taken is least.

In many ~~not~~ number of cases the real maximum rather than minimum. So in more general form "A ray of light travelling from one point to another by any number of reflections and refractions follows a path for which, compared with all other neighbouring paths, the time taken is either a minimum or max<sup>m</sup> or stationary."

Deduction of laws of reflection :-

Let a ray of light from a point A be reflected by a plane mirror  $zz'$  to another point B. Let  $i'$  and  $r'$  be the angle of




fig (ii)

H.  
Hip.  
HoO-Ray  
Thank You

AN APPLE A DAY ★  
KEEPS THE DOCTOR AWAY

NAVNEET

incident and reflection. Let  $z'$  &  $b$  be the length of  $\perp$  drawn from A and B on the mirror.  
 let  $zz' = a$ ,  $Oz = x$ ,  $Oz' = (c-x)$

The path  $AOB$  is travelled in air so the optical path between A and B is

$$I = OA + OB = \sqrt{a^2 + x^2} + \sqrt{b^2 + (c-x)^2} \quad \textcircled{1}$$

According to Fermat's principle O will have positions such that the optical path I is a min<sup>m</sup> or max<sup>m</sup>. So the first differential coefficient w.r.t  $x$  must be zero.

Now differentiating  $\textcircled{1}$  w.r.t  $x$  and equating the result to zero, we get

$$\frac{dI}{dx} = \frac{1}{2}(a^2 + x^2)^{-\frac{1}{2}}(2x) + \frac{1}{2}[b^2 + (c-x)^2]^{-\frac{1}{2}}(c-x)(-1) = 0$$

This can be expressed as

$$\frac{x}{\sqrt{a^2 + x^2}} = \frac{c-x}{\sqrt{b^2 + (c-x)^2}} \quad \textcircled{2}$$

from  $\textcircled{1}$  figure

$$\frac{x}{\sqrt{a^2 + x^2}} = \frac{zo}{AO} = \sin i$$

$$\text{and } \frac{c-x}{\sqrt{b^2 + (c-x)^2}} = \frac{oz'}{OB} = \sin r$$

from eq<sup>n</sup>  $\textcircled{2}$


$$\sin i = \sin r$$

$$\therefore i = r$$

which is law of reflection.

### Deduction of laws of Refraction :-

Let a ray of light from point A having RI  $i_1$  be refracted to another point B having RI  $i_2$  along the path  $AOB$ . Let 'i' &



$\gamma'$  be the angle of incidence and refraction. Let  $a'$  and  $b'$  be the lengths of  $\perp$  drawn from A and B to refracting surface. (3)

Let  $zz' = c$ ,  $Oz = x$ ,  $Oz' = (c-x)$

The optical path between A and B is given as

$$I = u_1(AO) + u_2(OB) \\ = u_1 \sqrt{a^2+x^2} + u_2 \sqrt{b^2+(c-x)^2} \quad \text{--- (3)}$$

Differentiating eq<sup>n</sup> (3) w.r.t  $x$  and equating the result to zero.

$$\frac{dI}{dx} = u_1 \left(\frac{1}{2}\right) (a^2+x^2)^{-\frac{1}{2}} (2x) + u_2 \left(\frac{1}{2}\right) [b^2+(c-x)^2]^{-\frac{1}{2}} (c-x)(-1) = 0$$

This can be expressed as

$$u_1 \frac{x}{\sqrt{a^2+x^2}} = u_2 \frac{c-x}{\sqrt{b^2+(c-x)^2}} \quad \text{--- (4)}$$

from fig (1)

$$\frac{ax}{\sqrt{a^2+x^2}} = \frac{Oz}{OA} = \sin i \quad \text{and}$$

$$\frac{c-x}{\sqrt{b^2+(c-x)^2}} = \frac{Oz'}{OB} = \sin r$$

$$\therefore u_1 \sin i = u_2 \sin r \quad (\text{from (4)})$$

$$\therefore \frac{\sin i}{\sin r} = \frac{u_2}{u_1} = n^2$$

This is known as Snell's law of refraction.

This is law of refraction: Further if incident ray, refracted ray and normal at the point of incidence all lies in same plane.

